



Maxforce[®]
FORTE

Cockroach Makhani or Cockroach Crispy?

We help you make sure that such items
don't exist on menus

Talk to Bayer Pest Management Experts



ACTIVE INGREDIENT

Fipronil 0.03% RB

RECOMMENDED USES

Effective against cockroaches like German cockroach, Brown-banded cockroach, American cockroach and Oriental cockroach commonly found in the food handling areas of households and commercial premises

MODE OF ACTION

Stomach poison

BENEFITS

- Baits are attractive to most species of cockroaches
- Highly effective at very low doses
- Target specific to cockroaches
- Ready to use - No measuring or mixing is required



Household Insecticide
for effective management
of cockroaches



APPLICATION RATE

Cockroach Species	Application Rate (100 mg drop = approx. 6 mm diameter)	
	Low Density Infestation ^a	High Density Infestation ^b
German Cockroach, Brown-banded Cockroach	1 drop/ m ²	2 drops/ m ²
American Cockroach, Oriental Cockroach	2 drops/ m ²	3 drops/ m ²

^a Low Density Infestation: Cockroaches rarely visible during the day
^b High Density Infestation: Cockroaches commonly visible during the day

ANTIDOTE

No specific antidote. Give symptomatic treatment

PACKS

Cartridge of 35 g

Corporate Office:

Crop Science - Environmental Science

Bayer House, Central Avenue, Hiranandani Estate
Thane (West) - 4000 607. Maharashtra, India.

Customer Care Cell:

E-mail: customercare@bayer.com

Tel.: 022 - 2531 1234

Toll Free: 1800 209 5666

Choose Safety Choose Bayer

www.environmentalscience.bayer.in

